

# Lecture Notes 14

---

## Connections

Suppose that we have a vector field  $X$  on a Riemannian manifold  $M$ . How can we measure how much  $X$  is changing at a point  $p \in M$  in the direction  $Y_p \in T_p M$ ? The main problem here is that there exists no canonical way to compare a vector in some tangent space of a manifold to a vector in another tangent space. Hence we need to impose a new kind of structure on a manifold. To gain some insight, we first study the case where  $M = \mathbf{R}^n$ .

### 0.1 Differentiation of vector fields in $\mathbf{R}^n$

Since each tangent space  $T_p \mathbf{R}^n$  is canonically isomorphic to  $\mathbf{R}^n$ , any vector field on  $\mathbf{R}^n$  may be identified as a mapping  $X: \mathbf{R}^n \rightarrow \mathbf{R}^n$ . Then for any  $Y_p \in T_p \mathbf{R}^n$  we define the *covariant derivative* of  $X$  with respect to  $Y_p$  as

$$\nabla_{Y_p} X := (Y_p(X^1), \dots, Y_p(X^n)).$$

Recall that  $Y_p(X^i)$  is the directional derivative of  $X^i$  at  $p$  in the direction of  $Y$ , i.e., if  $\gamma: (-\epsilon, \epsilon) \rightarrow M$  is any smooth curve with  $\gamma(0) = p$  and  $\gamma'(0) = Y$ , then

$$Y_p(X^i) = (X^i \circ \gamma)'(0) = \langle \text{grad } X^i(p), Y \rangle.$$

The last equality is an easy consequence of the chain rule. Now suppose that  $Y: \mathbf{R}^n \rightarrow \mathbf{R}^n$  is a vector field on  $\mathbf{R}^n$ ,  $p \xrightarrow{Y} Y_p$ , then we may define a new vector field on  $\mathbf{R}^n$  by

$$(\nabla_Y X)_p := \nabla_{Y_p} X.$$

Then the operation  $(X, Y) \mapsto \nabla_X Y$  may be thought of as a mapping  $\nabla: \mathcal{X}(\mathbf{R}^n) \times \mathcal{X}(\mathbf{R}^n) \rightarrow \mathcal{X}(\mathbf{R}^n)$ , where  $\mathcal{X}$  denotes the space of vector fields on  $\mathbf{R}^n$ .

Next note that if  $X \in \mathcal{X}(\mathbf{R}^n)$  is any vector field and  $f: M \rightarrow \mathbf{R}$  is a function, then we may define a new vector field  $fX \in \mathcal{X}(\mathbf{R}^n)$  by setting  $(fX)_p := f(p)X_p$  (do not confuse  $fX$ , which is a *vector field*, with  $Xf$  which is a *function* defined by  $Xf(p) := X_p(f)$ ). Now we observe that the covariant differentiation of vector fields on  $\mathbf{R}^n$  satisfies the following properties:

1.  $\nabla_Y(X_1 + X_2) = \nabla_Y X_1 + \nabla_Y X_2$

---

<sup>1</sup>Last revised: February 12, 2025

2.  $\nabla_Y(fX) = (Yf)\nabla_Y X + f\nabla_Y X$
3.  $\nabla_{Y_1+Y_2}X = \nabla_{Y_1}X + \nabla_{Y_2}X$
4.  $\nabla_{fY}X = f\nabla_Y X$

It is an easy exercise to check the above properties. Another good exercise to write down the pointwise versions of the above expressions. For instance note that item (2) implies that

$$\nabla_{Y_p}(fX) = (Y_p f)\nabla_{Y_p} X + f(p)\nabla_{Y_p} X,$$

for all  $p \in M$ .

## 0.2 Definition of connection and Christoffel symbols

Motivated by the Euclidean case, we define a connection  $\nabla$  on a manifold  $M$  as any mapping

$$\nabla: \mathcal{X}(M) \times \mathcal{X}(M) \rightarrow \mathcal{X}(M)$$

which satisfies the four properties mentioned above. We say that  $\nabla$  is smooth if whenever  $X$  and  $Y$  are smooth vector fields on  $M$ , then  $\nabla_Y X$  is a smooth vector field as well. Note that any manifold admits the trivial connection  $\nabla \equiv 0$ . In the next sections we study some nontrivial examples.

Here we describe how to express a connection in local charts. Let  $E_i$  be a basis for the tangent space of  $M$  in a neighborhood of a point  $p$ . For instance, choose a local chart  $(U, \phi)$  centered at  $p$  and set  $E_i(q) := d\phi_{\phi(q)}^{-1}(e_i)$  for all  $q \in U$ . Then if  $X$  and  $Y$  are any vector fields on  $M$ , we may write  $X = \sum_i X^i E_i$ , and  $Y = \sum_i Y^i E_i$  on  $U$ . Consequently, if  $\nabla$  is a connection on  $M$  we have

$$\nabla_Y X = \nabla_Y \left( \sum_i X^i E_i \right) = \sum_i (Y(X^i)E_i + X^i \nabla_Y E_i).$$

Now note that since  $(\nabla_{E_j} E_i)_p \in T_p M$ , for all  $p \in U$ , then it is a linear combination of the basis elements of  $T_p M$ . So we may write

$$\nabla_{E_j} E_i = \sum_k \Gamma_{ji}^k E_k$$

for some functions  $\Gamma_{ji}^k$  on  $U$  which are known as the *Christoffel symbols*. Thus

$$\begin{aligned} \nabla_Y X &= \sum_i \left( Y(X^i)E_i + X^i \sum_j \left( Y^j \sum_k \Gamma_{ji}^k E_k \right) \right) \\ &= \sum_k \left( Y(X^k) + \sum_{ij} Y^i X^j \Gamma_{ij}^k \right) E_k \end{aligned}$$

Conversely note that, a choice of the functions  $\Gamma_{ij}^k$  on any local neighborhood of  $M$  defines a connection on that neighborhood by the above expression. Thus we may define a connection on any manifold, by an arbitrary choice of Christoffel symbols in each local chart of some atlas of  $M$  and then using a partition of unity.

Next note that for every  $p \in U$  we have:

$$(\nabla_Y X)_p = \sum_k \left( Y_p(X^k) + \sum_{ij} Y^i(p) X^j(p) \Gamma_{ij}^k(p) \right) E_k(p). \quad (1)$$

This immediately shows that

**Theorem 0.1.** *For any point  $p \in M$ ,  $(\nabla_Y X)_p$  depends only on the value of  $Y$  at  $p$  and the restriction of  $X$  to any curve  $\gamma: (-\epsilon, \epsilon) \rightarrow M$  which belongs to the equivalence class of curves determined by  $Y_p$ .*  $\square$

Thus if  $p \in M$ ,  $Y_p \in T_p M$  and  $X$  is any vector field which is defined on an open neighborhood of  $p$ , then we may define

$$\nabla_{Y_p} X := (\nabla_Y X)_p$$

where  $Y$  is any extension of  $Y_p$  to a vector field in a neighborhood of  $p$ . Note that such an extension may always be found: for instance, if  $Y_p = \sum Y_p^i E_i(p)$ , where  $E_i$  are some local basis for tangent spaces in a neighborhood  $U$  of  $p$ , then we may set  $Y_q := \sum Y_p^i E_i(q)$  for all  $q \in U$ . By the previous proposition,  $(\nabla_Y X)_p$  does not depend on the choice of the local extension  $Y$ , so  $\nabla_{Y_p} X$  is well defined.

### 0.3 Induced connection on submanifolds

As we have already seen  $M$  admits a standard connection when  $M = \mathbf{R}^n$ . To give other examples of manifolds with a distinguished connection, we use the following observation.

**Lemma 0.2.** *Let  $\overline{M}$  be a manifold,  $M$  be an embedded submanifold of  $M$ , and  $X$  be a vector field of  $M$ . Then for every point  $p \in M$  there exists an open neighborhood  $\overline{U}$  of  $p$  in  $\overline{M}$  and a vector field  $\overline{X}$  defined on  $\overline{U}$  such that  $\overline{X}_p = X_p$  for all  $p \in M$ .*

*Proof.* Recall that, by the rank theorem, there exists a local chart  $(\overline{U}, \overline{\phi})$  of  $\overline{M}$  centered at  $p$  such that  $\overline{\phi}(U \cap M) = \mathbf{R}^{n-k}$  where  $k = \dim(\overline{M}) - \dim(M)$ . Now, note that  $d\overline{\phi}(X)$  is a vector field on  $\mathbf{R}^{n-k}$  and let  $Y$  be an extension of  $d\overline{\phi}(X)$  to  $\mathbf{R}^n$  (any vector field on a subspace of  $\mathbf{R}^n$  may be extended to all of  $\mathbf{R}^n$ ). Then set  $\overline{X} := d\overline{\phi}^{-1}(Y)$ .  $\square$

Now if  $\overline{M}$  is a Riemannian manifold with connection  $\overline{\nabla}$ , and  $M$  is any submanifold of  $\overline{M}$ , we may define a connection on  $M$  as follows. First note that for any  $p \in M$ ,

$$T_p \overline{M} = T_p M \oplus (T_p M)^\perp,$$

that is any vector  $X \in T_p\overline{M}$  may be written as sum of a vector  $X^\top \in T_pM$  (which is tangent to  $M$  and vector  $X^\perp := X - X^\top$  (which is normal to  $M$ ). So for any vector fields  $X$  and  $Y$  on  $M$  we define a new vector field on  $M$  by setting, for each  $p \in M$ ,

$$(\nabla_Y X)_p := (\overline{\nabla_{\overline{Y}} \overline{X}})_p^\top$$

where  $\overline{Y}$  and  $\overline{X}$  are local extensions of  $X$  and  $Y$  to vector fields on a neighborhood of  $p$  in  $M$ . Note  $(\nabla_Y X)_p$  is well-defined, because it is independent of the choice of local extensions  $\overline{X}$  and  $\overline{Y}$  by Theorem 0.1.

#### 0.4 Covariant derivative

We now describe how to differentiate a vector field along a curve in a manifold  $M$  with a connection  $\nabla$ . Let  $\gamma: I \rightarrow M$  be a smooth immersion, i.e.,  $d\gamma_t \neq 0$  for all  $t \in I$ , where  $I \subset \mathbf{R}$  is an open interval. By a vector field along  $\gamma$  we mean a mapping  $X: I \rightarrow TM$  such that  $X(t) \in T_{\gamma(t)}M$  for all  $t \in I$ . Let  $\mathcal{X}(\gamma)$  denote the space of vector fields along  $\gamma$ .

For any vector field  $X \in \mathcal{X}(\gamma)$ , we define another vector field  $D_\gamma X \in \mathcal{X}(\gamma)$ , called the covariant derivative of  $X$  along  $\gamma$ , as follows. First recall that  $\gamma$  is locally one-to-one by the inverse function theorem. Thus, by the previous lemma on the existence of local extensions of vector fields on embedded submanifolds, there exists an open neighborhood  $U$  of  $\gamma(t_0)$  and a vector field  $\overline{X}$  defined on  $U$  such that  $\overline{X}_{\gamma(t)} = X(t)$  for all  $t \in (t_0 - \epsilon, t_0 + \epsilon)$ . Set

$$D_\gamma X(t_0) := \nabla_{\gamma'(t_0)} \overline{X}.$$

Recall that  $\gamma'(t_0) := d\gamma_{t_0}(1) \in T_{\gamma(t_0)}M$ . By Theorem 0.1,  $D_\gamma X(t_0)$  is well defined, i.e., it does not depend on the choice of the local extension  $\overline{X}$ . Thus we obtain a mapping  $D_\gamma: \mathcal{X}(\gamma) \rightarrow \mathcal{X}(\gamma)$ . Note that if  $X, Y \in \mathcal{X}(\gamma)$ , then  $(X + Y)(t) := X(t) + Y(t) \in \mathcal{X}(\gamma)$ . Further, if  $f: I \rightarrow \mathbf{R}$  is any function then  $(fX)(t) := f(t)X(t) \in \mathcal{X}(\gamma)$ . It is easy to check that

$$D_\gamma(X + Y) = D_\gamma(X) + D_\gamma(Y) \quad \text{and} \quad D_\gamma(fX) = fD_\gamma(X).$$

**Proposition 0.3.** *If  $\gamma: I \rightarrow \mathbf{R}^n$ , and  $X \in \mathcal{X}(\gamma)$ , then  $D_\gamma X = X'$ . In particular,  $D_\gamma \gamma' = \gamma''$ .*

*Proof.* Let  $\overline{X}$  be a vector field on an open neighborhood of  $\gamma(t_0)$  such that

$$\overline{X}(\gamma(t)) = X(t),$$

for all  $t \in (t_0 - \epsilon, t_0 + \epsilon)$ . Then

$$D_\gamma X(t_0) = \nabla_{\gamma'(t_0)} \overline{X} = (\overline{X} \circ \gamma)'(t_0) = X'(t_0).$$

□

**Corollary 0.4.** *Let  $M$  be an immersed submanifold of  $\mathbf{R}^n$  with the induced connection  $\bar{\nabla}$ , and corresponding covariant derivative  $\bar{D}$ . Suppose  $\gamma: I \rightarrow M$  is an immersed curve, and  $X \in \mathcal{X}_M(\gamma)$  is a vector field along  $\gamma$  in  $M$ . Then  $\bar{D}_\gamma X = (X')^\top$ .  $\square$*

## 0.5 Geodesics

Note that, by the last exercise, the only curves  $\gamma: I \rightarrow \mathbf{R}^n$  with the property that

$$D_\gamma \gamma' \equiv 0$$

are given by  $\gamma(t) = at + b$ , which trace straight lines. With this motivation, we define a *geodesic* (which is meant to be a generalization of the concept of lines) as an immersed curve  $\gamma: I \rightarrow M$  which satisfies the above equality for all  $t \in I$ . A nice supply of examples of geodesics are provided by the following observation:

**Proposition 0.5.** *Let  $M \subset \mathbf{R}^n$  be an immersed submanifold, and  $\gamma: I \rightarrow M$  an immersed curve. Then  $\gamma$  is a geodesic of  $M$  (with respect to the induced connection from  $\mathbf{R}^n$ ) if and only if  $\gamma''^\top \equiv 0$ . In particular, if  $\gamma: I \rightarrow M$  is a geodesic, then  $\|\gamma'\| = \text{const}$ .  $\square$*

*Proof.* The first claim is an immediate consequence of the last two results. The last sentence follows from the Leibnitz rule for differentiating inner products in Euclidean space:  $\langle \gamma', \gamma' \rangle' = 2\langle \gamma'', \gamma' \rangle$ . Thus if  $\gamma''^\top \equiv 0$ , then  $\|\gamma'\|^2 = \text{const}$ .  $\square$

As an application of the last result, we can show that the geodesics on the sphere  $\mathbf{S}^2$  are those curves which trace a great circle with constant speed:

**Example 0.6** (Geodesics on  $\mathbf{S}^2$ ). A  $C^2$  immersion  $\gamma: I \rightarrow \mathbf{S}^2$  is a geodesic if and only if  $\gamma$  has constant speed and lies on a plane which passes through the center of the sphere, i.e., it traces a segment of a great circle.

First suppose that  $\gamma: I \rightarrow \mathbf{S}^2$  has constant speed, i.e.  $\|\gamma'\| = \text{const}$ ., and that  $\gamma$  traces a part of a great circle, i.e.,  $\langle \gamma, u \rangle = 0$  for some fixed vector  $u \in \mathbf{S}^2$  (which is the vector orthogonal to the plane in which  $\gamma$  lies). Since  $\langle \gamma', \gamma' \rangle = \|\gamma'\|^2$  is constant, it follows from the Leibnitz rule for differentiating the innerproduct that  $\langle \gamma'', \gamma' \rangle = 0$ . Furthermore, differentiating  $\langle \gamma, u \rangle = 0$  yields that  $\langle \gamma'', u \rangle = 0$ . So,  $\gamma''$  lies in the plane of  $\gamma$ , and is orthogonal to  $\gamma$ . So, since  $\gamma$  traces a circle,  $\gamma''$  must be parallel to  $\gamma$ . This in turn implies that  $\gamma''$  must be orthogonal to  $T_\gamma \mathbf{S}^2$ , since  $\gamma$  is orthogonal to  $T_\gamma \mathbf{S}^2$ . So we conclude that  $(\gamma'')^\top = 0$ .

Conversely, suppose that  $(\gamma'')^\top = 0$ . Then  $\gamma''$  is parallel to  $\gamma$ . So if  $u := \gamma \times \gamma'$ , then  $u' = \gamma' \times \gamma' + \gamma \times \gamma'' = 0 + 0 = 0$ . So  $u$  is constant. But  $\gamma$  is orthogonal to  $u$ , so  $\gamma$  lies in the plane which passes through the origin and is orthogonal to  $u$ . Finally,  $\gamma$  has constant speed by the last proposition.

## 0.6 Ordinary differential equations

In order to prove an existence and uniqueness result for geodesic in the next section we need to develop first a basic result about differential equations:

**Theorem 0.7.** *Let  $U \subset \mathbf{R}^n$  be an open set and  $F: U \rightarrow \mathbf{R}^n$  be  $C^1$ , then for every  $x_0 \in U$ , there exists an  $\bar{\epsilon} > 0$  such that for every  $0 < \epsilon < \bar{\epsilon}$  there exists a unique curve  $x: (-\epsilon, \epsilon) \rightarrow U$  with  $x(0) = x_0$  and  $x'(t) = F(x(t))$ .*

Note that, from the geometric point of view the above theorem states that there passes an integral curve through every point of a vector field. To prove this result we need a number of preliminary results. Let  $I \subset \mathbf{R}$  be an interval,  $(X, d)$  be a compact metric space, and  $\Gamma(I, X)$  be the space of maps  $\gamma: I \rightarrow X$ . For every pair of curves  $\gamma_1, \gamma_2 \in \Gamma(I, X)$  set

$$\delta(\gamma_1, \gamma_2) := \sup_{t \in I} d(\gamma_1(t), \gamma_2(t)).$$

It is easy to check that  $(\Gamma, \delta)$  is a metric space. Now let  $C(I, X) \subset \Gamma(I, X)$  be the subspace of consisting of *continuous* curves.

**Lemma 0.8.**  *$(C, \delta)$  is a complete metric space.*

*Proof.* Let  $\gamma_i \in C$  be a Cauchy sequence. Then, for every  $t \in I$ ,  $\gamma_i(t)$  is a Cauchy sequence in  $X$ . So  $\gamma_i(t)$  converges to a point  $\bar{\gamma}(t) \in X$  (since every compact metric space is complete). Thus we obtain a mapping  $\bar{\gamma}: I \rightarrow X$ . We claim that  $\bar{\gamma}$  is continuous which would complete the proof. By the triangular inequality,

$$\begin{aligned} d(\bar{\gamma}(s), \bar{\gamma}(t)) &\leq d(\bar{\gamma}(s), \gamma_i(s)) + d(\gamma_i(s), \gamma_i(t)) + d(\gamma_i(t), \bar{\gamma}(t)) \\ &\leq 2\delta(\bar{\gamma}, \gamma_i) + d(\gamma_i(s), \gamma_i(t)). \end{aligned}$$

So, since  $\gamma_i$  is continuous,

$$\lim_{t \rightarrow s} d(\bar{\gamma}(s), \bar{\gamma}(t)) \leq 2\delta(\bar{\gamma}, \gamma_i).$$

All we need then is to check that  $\lim_{i \rightarrow \infty} \delta(\bar{\gamma}, \gamma_i) = 0$ : Given  $\epsilon > 0$ , choose  $i$  sufficiently large so that  $\delta(\gamma_i, \gamma_j) < \epsilon$  for all  $j \geq i$ . Then, for all  $t \in I$ ,  $d(\gamma_i(t), \gamma_j(t)) \leq \epsilon$ , which in turn yields that  $d(\gamma_i(t), \bar{\gamma}(t)) \leq \epsilon$ . So  $\delta(\gamma_i, \bar{\gamma}) \leq \epsilon$ .  $\square$

Now we are ready to prove the main result of this section:

*Proof of Theorem 0.7.* Let  $B = B_r^n(x_0)$  denote a ball of radius  $r$  centered at  $x_0$ . Choose  $r > 0$  so small that that  $\bar{B} \subset U$ . For any continuous curve  $\alpha \in C((-\epsilon, \epsilon), \bar{B})$  we may define another continuous curve  $s(\alpha) \in ((-\epsilon, \epsilon), \mathbf{R}^n)$  by

$$s(\alpha)(t) := x_0 + \int_0^t F(a(u)) du.$$

We claim that if  $\epsilon$  is small enough, then  $s(\alpha) \in C((-\epsilon, \epsilon), \overline{B})$ . To see this note that

$$\|s(\alpha)(t) - x_0\| = \left\| \int_0^t F(\alpha(u)) du \right\| \leq \int_0^t \|F(\alpha(u))\| du \leq \epsilon \sup_{\overline{B}} \|F\|.$$

So setting  $\epsilon \leq r / \sup_{\overline{B}} \|F\|$ , we may then assume that

$$s: C((-\epsilon, \epsilon), \overline{B}) \rightarrow C((-\epsilon, \epsilon), \overline{B}).$$

Next note that for every  $\alpha, \beta \in C((-\epsilon, \epsilon), \overline{B})$ , we have

$$\delta(s(\alpha), s(\beta)) = \sup_t \left\| \int_0^t F(\alpha(u)) - F(\beta(u)) du \right\| \leq \sup_t \int_0^t \|F(\alpha(u)) - F(\beta(u))\| du.$$

Further recall that, since  $F$  is  $C^1$ , by the mean value theorem there is a constant  $K$  such that

$$\|F(x) - F(y)\| \leq K \|x - y\|,$$

for all  $x, y \in \overline{B}$  (in particular recall that we may set  $K := \sqrt{n} \sup_{\overline{B}} |D_j F^i|$ ). Thus

$$\int_0^t \|F(\alpha(u)) - F(\beta(u))\| du \leq K \int_0^t \|\alpha(u) - \beta(u)\| du \leq K \epsilon \delta(\alpha, \beta).$$

So we conclude that

$$\delta(s(\alpha), s(\beta)) \leq K \epsilon \delta(\alpha, \beta).$$

Now assume that  $\epsilon < 1/K$  (in addition to the earlier assumption that  $\epsilon \leq r / \sup_{\overline{B}} \|F\|$ ), then,  $s$  must have a unique fixed point since it is a contraction mapping. So for every  $0 < \epsilon < \bar{\epsilon}$  where

$$\bar{\epsilon} := \min \left\{ \frac{r}{\sup_{\overline{B}} \|F\|}, \frac{1}{\sqrt{n} \sup_{\overline{B}} |D_j F^i|} \right\}$$

there exists a unique curve  $x: (-\epsilon, \epsilon) \rightarrow \overline{B}$  such that  $x(0) = s(x)(0) = x_0$ , and  $x'(t) = s(x)'(t) = F(x(t))$ .

It only remains to show that  $x: (-\epsilon, \epsilon) \rightarrow U$  is also the unique curve with  $x(0) = x_0$  and  $x'(t) = F(x(t))$ , i.e., we have to show that if  $y: (-\epsilon, \epsilon) \rightarrow U$  is any curve with  $y(0) = x_0$  and  $y'(t) = F(y(t))$ , then  $y = x$  (so far we have proved this only for  $y: (-\epsilon, \epsilon) \rightarrow \overline{B}$ ). To see this recall that  $\epsilon \leq r / \sup_{\overline{B}} \|F\|$  where  $r$  is the radius of  $\overline{B}$ . Thus

$$\|y(t) - x_0\| \leq \int_0^t \|y'(u)\| du = \int_0^t \|F(y(u))\| du \leq \epsilon \sup_{\overline{B}} \|F\| \leq r.$$

So the image of  $y$  lies in  $\overline{B}$ , and therefore we must have  $y = x$ . □

## 0.7 Existence and uniqueness of geodesics

Note that for every point  $p \in \mathbf{R}^n$  and vector  $X \in T_p \mathbf{R}^n \simeq \mathbf{R}^n$ , we may find a geodesic through  $p$  and with velocity vector  $X$  at  $p$ , which is given simply by  $\gamma(t) = p + Xt$ . Here we show that all manifolds with a connection share this property:

**Theorem 0.9.** *Let  $M$  be a manifold with a connection. Then for every  $p \in M$  and  $X \in T_p M$  there exists an  $\bar{\epsilon} > 0$  such that for every  $0 < \epsilon < \bar{\epsilon}$  there is a unique geodesic  $\gamma: (-\epsilon, \epsilon) \rightarrow M$  with  $\gamma(0) = p$  and  $\gamma'(0) = X$ .*

To prove this theorem, we need to record some preliminary observations. Let  $M$  and  $\widetilde{M}$  be manifolds with connections  $\nabla$  and  $\widetilde{\nabla}$  respectively. We say that a diffeomorphism  $f: M \rightarrow \widetilde{M}$  is connection preserving provided that

$$(\nabla_Y X)_p = (\widetilde{\nabla}_{df(Y)} df(X))_{f(p)}$$

for all  $p \in M$  and all vector fields  $X, Y \in \mathcal{X}(M)$ . It is an immediate consequence of the definitions that

**Lemma 0.10.** *Let  $f: M \rightarrow \widetilde{M}$  be a connection preserving diffeomorphism. Then  $\gamma: I \rightarrow M$  is a geodesic if and only if  $f \circ \gamma$  is a geodesic.*  $\square$

Note that if  $f: M \rightarrow \widetilde{M}$  is a diffeomorphism, and  $M$  has a connection  $\nabla$ , then  $f$  induces a connection  $\widetilde{\nabla}$  on  $\widetilde{M}$  by

$$(\widetilde{\nabla}_{\widetilde{Y}} \widetilde{X})_{\widetilde{p}} := (\nabla_{df^{-1}(\widetilde{X})} df^{-1}(\widetilde{Y}))_{f^{-1}(\widetilde{p})}.$$

It is clear that then  $f: M \rightarrow \widetilde{M}$  will be connection preserving. So we may conclude that

**Lemma 0.11.** *Let  $(U, \phi)$  be a local chart of  $M$ , then  $\gamma: I \rightarrow U$  is a geodesic if and only if  $\phi \circ \gamma$  is a geodesic with respect to the connection induced on  $\mathbf{R}^n$  by  $\phi$ .*  $\square$

Now we are ready to prove the main result of this section:

*Proof of Theorem 0.9.* Let  $(U, \phi)$  be a local chart of  $M$  centered at  $p$  and let  $\nabla$  be the connection which is induced on  $\phi(U) = \mathbf{R}^n$  by  $\phi$ . We will show that there exists an  $\bar{\epsilon} > 0$  such that for every  $0 < \epsilon < \bar{\epsilon}$  there is a unique geodesic  $c: (-\epsilon, \epsilon) \rightarrow \mathbf{R}^n$ , with respect to the induced connection, which satisfies the initial conditions

$$c(0) = \phi(p) \quad \text{and} \quad c'(0) = d\phi_p(X).$$

Then, by a previous lemma,  $\gamma := \phi^{-1} \circ c: (-\epsilon, \epsilon) \rightarrow M$  will be a geodesic on  $M$  with  $\gamma(0) = p$  and  $\gamma'(0) = X$ . Furthermore,  $\gamma$  will be unique. To see this suppose that  $\bar{\gamma}: (-\epsilon, \epsilon) \rightarrow M$  is another geodesic with  $\bar{\gamma}(0) = p$  and  $\bar{\gamma}'(0) = X$ . Let  $\epsilon'$  be the supremum of  $t \in [0, \epsilon]$  such that  $\bar{\gamma}(-t, t) \subset U$ , and set  $\bar{c} := \phi \circ \bar{\gamma}|_{(-\epsilon', \epsilon')}$ .



Then, by Theorem 0.7,  $c = \bar{c}$  on  $(-\epsilon', \epsilon')$ , because  $\epsilon' < \bar{\epsilon}$ . So it follows that  $\gamma = \bar{\gamma}$  on  $(-\epsilon', \epsilon')$ , and we are done if  $(-\epsilon', \epsilon') = (-\epsilon, \epsilon)$ . This is indeed the case, for otherwise,  $(-\epsilon' - \delta, \epsilon' + \delta) \subset (-\epsilon, \epsilon)$ , for some  $\delta > 0$ . Further  $\bar{\gamma}(\pm\epsilon') = \gamma(\pm\epsilon') \in U$ . So if  $\delta$  is sufficiently small, then  $\bar{\gamma}(-\epsilon' - \delta, \epsilon' + \delta) \subset U$ , which contradicts the definition of  $\epsilon'$ .

So all we need is to establish the existence and uniqueness of the geodesic  $c: (-\epsilon, \epsilon) \rightarrow \mathbf{R}^n$  mentioned above. For  $c$  to be a geodesic we must have

$$D_c c' \equiv 0.$$

We will show that this may be written as a system of ordinary differential equations. To see this first recall that

$$D_c \dot{c}(t) = \nabla_{\dot{c}(t)} \bar{c}$$

where  $\bar{c}$  is a vector field in a neighborhood of  $c(t)$  which is a local extension of  $\dot{c}$ , i.e.,

$$\bar{c}(c(t)) = \dot{c}(t).$$

By (1) we have

$$\nabla_{\dot{c}(t)} \bar{c} = \sum_k \left( \dot{c}(t)(\bar{c}^k) + \sum_{ij} \dot{c}^i(t) \dot{c}^j(t) \Gamma_{ij}^k(c(t)) \right) e_k,$$

where  $e_i$  are the standard basis of  $\mathbf{R}^n$  and  $\Gamma_{ij}^k(p) = \langle (\nabla_{e_i} e_j)_p, e_k \rangle$ . But

$$\dot{c}(t)(\bar{c}^k) = (\bar{c}^k \circ c)'(t) = (\dot{c}^k)'(t) = \ddot{c}^k(t).$$

So  $D_c c' \equiv 0$  if and only if

$$\ddot{c}^k(t) + \sum_{ij} \dot{c}^i(t) \dot{c}^j(t) \Gamma_{ij}^k(c(t)) = 0$$

for all  $t \in I$  and all  $k$ . This is a system of  $n$  second order ordinary differential equations (ODEs), which we may rewrite as a system of  $2n$  first order ODEs, via substitution  $\dot{c} = v$ . Then we have

$$\begin{aligned} \dot{c}^k(t) &= v^k(t) \\ \dot{v}^k(t) &= - \sum_{ij} v^i(t) v^j(t) \Gamma_{ij}^k(c(t)). \end{aligned}$$

Now let  $\alpha(t) := (c(t), v(t))$ , and define  $F: \mathbf{R}^{2n} \rightarrow \mathbf{R}^{2n}$ ,  $F = (F^1, \dots, F^{2n})$  by

$$F^\ell(x, y) = y_\ell, \quad \text{and} \quad F^{\ell+n}(x, y) = - \sum_{ij} y^i y^j \Gamma_{ij}^\ell(x)$$

for  $\ell = 1, \dots, n$ . Then the system of  $2n$  ODEs mentioned above may be rewritten as

$$\alpha'(t) = F(\alpha(t)),$$

which has a unique solution with initial conditions  $\alpha(0) = (\phi(p), d\phi(X))$ . □

## 0.8 Parallel translation

Let  $M$  be a manifold with a connection, and  $\gamma: I \rightarrow M$  be an immersed curve. Then we say that a vector field  $X \in \mathcal{X}(\gamma)$  is *parallel* along  $\gamma$  if

$$D_\gamma X \equiv 0.$$

Thus, in this terminology,  $\gamma$  is a geodesic if its velocity vector field is parallel. Further note that if  $M$  is a submanifold of  $\mathbf{R}^n$ , then, by the earlier results in this section,  $X$  is parallel along  $\gamma$  if and only if  $(X')^\top \equiv 0$ .

**Example 0.12.** Let  $M$  be a two dimensional manifold immersed in  $\mathbf{R}^n$ ,  $\gamma: I \rightarrow M$  be a geodesic of  $M$ , and  $X \in \mathcal{X}_M(\gamma)$  be a vector field along  $\gamma$  in  $M$ . Then  $X$  is parallel along  $\gamma$  if and only if  $X$  has constant length and the angle between  $X(t)$  and  $\gamma'(t)$  is constant as well. To see this note that  $(\gamma'')^\top \equiv 0$  since  $\gamma$  is a geodesic; therefore,

$$\langle X, \gamma' \rangle' = \langle X', \gamma' \rangle + \langle X, \gamma'' \rangle = \langle X', \gamma' \rangle.$$

So, if  $(X')^\top = 0$ , then it follows that  $\langle X, \gamma' \rangle$  is constant which since  $\gamma'$  and  $X$  have both constant lengths, implies that the angle between  $X$  and  $\gamma'$  is constant. Conversely, suppose that  $X$  has constant length and makes a constant angle with  $\gamma'$ . Then  $\langle X, \gamma' \rangle$  is constant, and the displayed expression above implies that  $\langle X, \gamma' \rangle = 0$  is constant. Furthermore,  $0 = \langle X, X' \rangle = 2\langle X, X' \rangle$ . So  $X'(t)$  is orthogonal to both  $X(t)$  and  $\gamma'(t)$ . If  $X(t)$  and  $\gamma'(t)$  are linearly dependent, then this implies that  $X'(t)$  is orthogonal to  $T_{\gamma(t)}M$ , i.e.,  $(X')^\top \equiv 0$ . If  $X(t)$  and  $\gamma'(t)$  are linearly independent, then  $(X')^\top = D_\gamma(X) = D_\gamma(f\gamma') = fD_\gamma(\gamma') \equiv 0$ .

**Example 0.13** (Foucault's Pendulum). Here we explicitly compute the parallel translation of a vector along a meridian of the sphere. To this end let

$$X(\theta, \phi) := (\cos(\theta) \sin(\phi), \sin(\theta) \sin(\phi), \cos(\phi))$$

be the standard parametrization or local coordinates for  $\mathbf{S}^2 - \{(0, 0, \pm 1)\}$ . Suppose that we want to parallel transport a given unit vector  $V_0 \in T_{X(\theta_0, \phi_0)}\mathbf{S}^2$  along the meridian  $X(\theta, \phi_0)$ , where we identify tangent space of  $\mathbf{S}^2$  with subspaces of  $\mathbf{R}^3$ . So we need to find a mapping  $V: [0, 2\pi] \rightarrow \mathbf{S}^2$  such that  $V(0) = V_0$  and  $V'(\theta) \perp T_{X(\theta, \phi_0)}\mathbf{S}^2$ . The latter condition is equivalent to the requirement that

$$V'(\theta) = \lambda(\theta)X(\theta, \phi_0), \tag{2}$$

since the normal to  $\mathbf{S}^2$  at the point  $X(\theta, \phi)$  is just  $X(\theta, \phi)$  itself. To solve the above differential equation, let

$$E_1(\theta) := \frac{\partial X / \partial \theta(\theta, \phi_0)}{\|\partial X / \partial \theta(\theta, \phi_0)\|} = (-\sin(\theta), \cos(\theta), 0),$$

and

$$E_2(\theta) := \frac{\partial X / \partial \phi(\theta, \phi_0)}{\|\partial X / \partial \phi(\theta, \phi_0)\|} = (\cos(\theta) \cos(\phi_0), \sin(\theta) \cos(\phi_0), -\sin(\phi_0)).$$

Now note that  $\{E_1(\theta), E_2(\theta)\}$  forms an orthonormal basis for  $T_{X(\theta_0, \phi_0)}\mathbf{S}^2$ . Thus (2) is equivalent to

$$\langle V'(\theta), E_1(\theta) \rangle = 0 \quad \text{and} \quad \langle V'(\theta), E_2(\theta) \rangle = 0. \quad (3)$$

So it remains to solve this differential equation. To this end first recall that since  $V_0$  has unit length, and parallel translation preserves length, we may write

$$V(\theta) = \cos(\alpha(\theta))E_1(\theta) + \sin(\alpha(\theta))E_2(\theta).$$

So differentiation yields that

$$V' = E_1' \cos(\alpha) - \sin(\alpha)\alpha' E_1 + \sin(\alpha)E_2' + \cos(\alpha)\alpha' E_2.$$

Further, it is easy to compute that

$$E_1' = -\cos(\phi_0)E_2 - \sin(\phi_0)E_3 \quad \text{and} \quad E_2' = \cos(\phi_0)E_1,$$

where  $E_3(\theta) := X(\theta, \phi_0)$ . Thus we obtain:

$$V' = \sin(\alpha)(\cos(\phi_0) - \alpha')E_1 + \cos(\alpha)(\alpha' - \cos(\phi_0))E_2 + (*)E_3.$$

So for (3) to be satisfied, we must have  $\alpha' = \cos(\phi_0)$  or

$$\alpha(\theta) = \cos(\phi_0)t + \alpha(0),$$

which in turns determines  $V$ . Note in particular that the total rotation of  $V$  with respect to the meridian  $X(\theta, \phi_0)$  is given by

$$\alpha(2\pi) - \alpha(0) = \int_0^{2\pi} \alpha' d\theta = 2\pi \cos(\phi_0).$$

Thus

$$\phi_0 = \cos^{-1} \left( \frac{\alpha(2\pi) - \alpha(0)}{2\pi} \right).$$

The last equation gives the relation between the precession of the swing plane of a pendulum during a 24 hour period, and the longitude of the location of that pendulum on earth, as first observed by the French Physicist Leon Foucault in 1851.

**Lemma 0.14.** *Let  $I \subset \mathbf{R}$  and  $U \subset \mathbf{R}^n$  be open subsets and  $F: I \times U \rightarrow \mathbf{R}^n$ , be  $C^1$ . Then for every  $t_0 \in I$  and  $x_0 \in U$  there exists an  $\bar{\epsilon} >$  such that for every  $0 < \epsilon < \bar{\epsilon}$  there is a unique curve  $x: (t_0 - \epsilon, t_0 + \epsilon) \rightarrow \mathbf{R}^n$  with  $x(t_0) = x_0$  and  $x'(t) = F(t, x(t))$ .*

*Proof.* Define  $\bar{F}: I \times U \rightarrow \mathbf{R}^{n+1}$  by  $\bar{F}(t, x) := (1, F(t, x))$ . Then, by Theorem 0.7, there exists an  $\bar{\epsilon} > 0$  and a unique curve  $\bar{x}: (t_0 - \epsilon, t_0 + \epsilon) \rightarrow \mathbf{R}^{n+1}$ , for every  $0 < \epsilon < \bar{\epsilon}$ , such that  $\bar{x}(t_0) = (1, x_0)$  and  $\bar{x}'(t) = \bar{F}(\bar{x}(t))$ . It follows then that  $\bar{x}(t) = (1, x(t))$ , for some unique curve  $x: (t_0 - \epsilon, t_0 + \epsilon) \rightarrow \mathbf{R}^n$ . Thus  $\bar{F}(\bar{x}(t)) = (1, F(t, x(t)))$ , and it follows that  $x'(t) = F(t, x(t))$ .  $\square$

**Lemma 0.15.** *Let  $A(t)$ ,  $t \in I$ , be a  $C^1$  one-parameter family of matrices. Then for every  $x_0 \in \mathbf{R}^n$  and  $t_0 \in I$ , there exists a unique curve  $x: I \rightarrow \mathbf{R}^n$  with  $x(t_0) = x_0$  such that  $x'(t) = A(t) \cdot x(t)$ .*

*Proof.* Define  $F: I \times \mathbf{R}^n \rightarrow \mathbf{R}^n$  by  $F_t(x) = A(t) \cdot x$ . By the previous lemma, there exists a unique curve  $x: (t_0 - \epsilon, t_0 + \epsilon) \rightarrow \mathbf{R}^n$  with  $x(t_0) = x_0$  such that  $F_t(x(t)) = x'(t)$  for all  $t \in (t_0 - \epsilon, t_0 + \epsilon)$ .

Now let  $J \subset I$  be the union of all open intervals in  $I$  which contains  $t_0$  and such that  $x'(t) = F(x(t))$  for all  $t$  in those intervals. Then  $J$  is open in  $I$  and nonempty. All we need then is to show that  $J$  is closed, for then it would follow that  $J = I$ . Suppose that  $\bar{t}$  is a limit point of  $J$  in  $I$ . Just as we argued in the first paragraph, there exists a curve  $y: (\bar{t} - \bar{\epsilon}, \bar{t} + \bar{\epsilon}) \rightarrow \mathbf{R}^n$  such that  $y'(t) = F(y(t))$  and  $y'(\bar{t}) \neq 0$ . Thus we may assume that  $y' \neq 0$  on  $(\bar{t} - \bar{\epsilon}, \bar{t} + \bar{\epsilon})$ , after replacing  $\bar{\epsilon}$  by a smaller number. In particular  $y'(\bar{t}) \neq 0$  for some  $\tilde{t} \in (\bar{t} - \bar{\epsilon}, \bar{t} + \bar{\epsilon}) \cap J$ , and there exists a matrix  $B$  such that  $B \cdot y'(\tilde{t}) = x'(\tilde{t})$ .

Now let  $\bar{y}(t) := B \cdot y(t)$ . Since  $F(y(t)) = y'(t)$ , we have  $F(\bar{y}(t)) = \bar{y}'(t)$ . Further, by construction  $\bar{y}(\tilde{t}) = x(\tilde{t})$ , so by uniqueness part of the previous result we must have  $\bar{y} = x$  on  $(\bar{t} - \bar{\epsilon}, \bar{t} + \bar{\epsilon}) \cap J$ . Thus  $x$  is defined on  $J \cup (\bar{t} - \bar{\epsilon}, \bar{t} + \bar{\epsilon})$ . But  $J$  was assumed to be maximal. So  $(\bar{t} - \bar{\epsilon}, \bar{t} + \bar{\epsilon}) \subset J$ . In particular  $\bar{t} \in J$ , which completes the proof that  $J$  is closed in  $I$ .  $\square$

**Theorem 0.16.** *Let  $X: I \rightarrow M$  be a  $C^1$  immersion. For every  $t_0 \in I$  and  $X_0 \in T_{\gamma(t_0)}M$ , there exists a unique parallel vector field  $X \in \mathcal{X}(\gamma)$  such that  $X(t_0) = X_0$ .*

*Proof.* First suppose that there exists a local chart  $(U, \phi)$  such that  $\gamma: I \rightarrow U$  is an embedding. Let  $\bar{X}$  be a vector field on  $U$  and set  $X(t) := \bar{X}(\gamma(t))$ . By (1),

$$D_\gamma(X)(t) = \nabla_{\gamma'(t)} \bar{X} = \sum_k \left( \gamma'(t)(\bar{X}^k) + \sum_{ij} \gamma^i(t) X^j(t) \Gamma_{ij}^k(\gamma(t)) \right) E_k(\gamma(t)).$$

Further note that

$$\gamma'(t) \bar{X} = (\bar{X} \circ \gamma)'(t) = X'(t).$$

So, in order for  $X$  to be parallel along  $\gamma$  we need to have

$$\dot{X}^k + \sum_{ij} \gamma^i(t) \Gamma_{ij}^k(\gamma(t)) X^j(t) = 0,$$

for  $k = 1, \dots, n$ . This is a linear system of ODE's in terms of  $X^i$ , and therefore by the previous lemma it has a unique solution on  $I$  satisfying the initial conditions  $X^i(t_0) = X_0^i$ .

Now let  $J \subset I$  be a compact interval which contains  $t_0$ . There exists a finite number of local charts of  $M$  which cover  $\gamma(J)$ . Consequently there exist subintervals  $J_1, \dots, J_n$  of  $J$  such that  $\gamma$  embeds each  $J_i$  into a local chart of  $M$ . Suppose that  $t_0 \in J_\ell$ , then, by the previous paragraph, we may extend  $X_0$  to a parallel vector field defined on  $J_\ell$ . Take an element of this extension which lies in a subinterval  $J_{\ell'}$  intersecting  $J_\ell$  and apply the previous paragraph to  $J_{\ell'}$ . Repeating this procedure, we obtain a parallel vector field on each  $J_i$ . By the uniqueness of each local extension mentioned above, these vector fields coincide on the overlaps of  $J_i$ . Thus we obtain a well-defined vector field  $X$  on  $J$  which is a parallel extension of  $X_0$ . Note that if  $\bar{J}$  is any other compact subinterval of  $I$  which contains  $t_0$ , and  $\bar{X}$  is the parallel extension of  $X_0$  on  $\bar{J}$ , then  $X$  and  $\bar{X}$  coincide on  $J \cap \bar{J}$ , by the uniqueness of local parallel extensions. Thus, since each point of  $I$  is contained in a compact subinterval containing  $t_0$ , we may consistently define  $X$  on all of  $I$ .

Finally let  $\bar{X}$  be another parallel extension of  $X_0$  defined on  $I$ . Let  $A \subset I$  be the set of points where  $\bar{X} = X$ . Then  $A$  is closed, by continuity of  $\bar{X}$  and  $X$ . Further  $A$  is open by the uniqueness of local extensions. Furthermore,  $A$  is nonempty since  $t_0 \in A$ . So  $A = I$  and we conclude that  $X$  is unique.  $\square$

Using the previous result we now define, for every  $X_0 \in T_{\gamma(t_0)}M$ ,

$$P_{\gamma, t_0, t}(X_0) := X(t)$$

as the *parallel transport* of  $X_0$  along  $\gamma$  to  $T_{\gamma(t)}M$ . Thus we obtain a mapping from  $T_{\gamma(t_0)}M$  to  $T_{\gamma(t)}M$ .

**Exercise 0.17.** Show that  $P_{\gamma, t_0, t}: T_{\gamma(t_0)}M \rightarrow T_{\gamma(t)}M$  is an isomorphism (*Hint:* Use the fact that  $D_\gamma: \mathcal{X}(\gamma) \rightarrow \mathcal{X}(\gamma)$  is linear). Also show that  $P_{\gamma, t_0, t}$  depends on the choice of  $\gamma$ .

**Exercise 0.18.** Show that

$$\nabla_{\gamma'(t_0)} X = \lim_{t \rightarrow t_0} \frac{P_{\gamma, t_0, t}^{-1}(X_{\gamma(t)}) - X_{\gamma(t_0)}}{t - t_0}.$$

(*Hint:* Use a parallel frame along  $\gamma$ .)