

h -PRINCIPLES FOR SMOOTH CURVES AND KNOTS WITH PRESCRIBED CURVATURE

MOHAMMAD GHOMI AND MATTEO RAFFAELLI

ABSTRACT. We show that smooth curves with prescribed curvature satisfy a \mathcal{C}^1 -dense h -principle in the space of immersed curves in Euclidean space. More precisely, every $\mathcal{C}^{\alpha \geq 2}$ curve with nonvanishing curvature in $\mathbf{R}^{n \geq 3}$ can be \mathcal{C}^1 -approximated by \mathcal{C}^α curves of any larger curvature, prescribed as a function of arclength. It follows that there exist \mathcal{C}^∞ knots of prescribed curvature in every isotopy class of closed curves embedded in \mathbf{R}^3 .

1. INTRODUCTION

Wasem [13] showed that every \mathcal{C}^2 curve in Euclidean space $\mathbf{R}^{n \geq 3}$ may be \mathcal{C}^1 -approximated by curves of larger prescribed curvature. The proof, which is based on the work of Nash and Kuiper [1, 9, 11], requires the target curvature function to be \mathcal{C}^∞ , but produces only \mathcal{C}^2 curves. Adopting a more geometric approach, we refine Wasem's result to obtain optimal regularity. We also achieve tighter control over the speed and position of the approximating curves.

To state our main result, let Γ be the interval $[a, b] \subset \mathbf{R}$ or a topological circle $\mathbf{R}/((b-a)\mathbf{Z})$, and $\mathcal{C}^\alpha(\Gamma, \mathbf{R}^n)$ be the space of \mathcal{C}^α maps $f: \Gamma \rightarrow \mathbf{R}^n$ with the topology induced by the \mathcal{C}^α -norm $|\cdot|_\alpha$. The space of (immersed) *curves* $\text{Imm}^{\alpha \geq 1}(\Gamma, \mathbf{R}^n) \subset \mathcal{C}^\alpha(\Gamma, \mathbf{R}^n)$ consists of maps with nonvanishing derivative. The *curvature* of $f \in \text{Imm}^2(\Gamma, \mathbf{R}^n)$ is given by $\kappa := |T'|/|f'|$, where $T := f'/|f'|$ is the *tantrix* of f .

Theorem 1.1. *Let $f \in \text{Imm}^{\alpha \geq 2}(\Gamma, \mathbf{R}^{n \geq 3})$ be a curve with curvature $\kappa > 0$, and $\tilde{\kappa}: \Gamma \rightarrow \mathbf{R}$ be a $\mathcal{C}^{\alpha-2}$ function with $\tilde{\kappa} > \kappa$. Then, for any $\varepsilon > 0$, there exists a curve $\tilde{f} \in \text{Imm}^\alpha(\Gamma, \mathbf{R}^n)$ with curvature $\tilde{\kappa}$ and $|\tilde{f} - f|_1 \leq \varepsilon$. If f has unit speed, then so does \tilde{f} . Furthermore, \tilde{f} can be made tangent to f at any finite set of points prescribed along Γ .*

The last two properties of \tilde{f} are additional features of our method, not granted by the Nash–Kuiper approach in [13]. If ε in Theorem 1.1 is sufficiently small, $h_t := (1-t)f + t\tilde{f}$, $t \in [0, 1]$, is a homotopy in $\text{Imm}^\alpha(\Gamma, \mathbf{R}^n)$. Thus, in the terminology of Gromov [7] or Eliashberg [2], the above result establishes a \mathcal{C}^1 -dense h -principle for smooth curves of prescribed curvature. Similar h -principles for curves of constant curvature or constant torsion were obtained in [4–6]. Theorem 1.1 has the following quick application. Let $\text{Emb}^\alpha(\Gamma, \mathbf{R}^3) \subset \text{Imm}^\alpha(\Gamma, \mathbf{R}^3)$ be the space of injective curves, which are called *knots* when Γ is a circle.

Corollary 1.2. *Let $f \in \text{Emb}^{\alpha \geq 2}(\Gamma, \mathbf{R}^3)$ be a knot, and $\tilde{\kappa}: \Gamma \rightarrow \mathbf{R}$ be a positive $\mathcal{C}^{\alpha-2}$ function. Then f is isotopic in $\text{Emb}^\alpha(\Gamma, \mathbf{R}^3)$ to a constant-speed knot of curvature $\tilde{\kappa}$.*

Date: Last revised on May 18, 2026.

1991 Mathematics Subject Classification. Primary 53A04, 57K10; Secondary 58C35, 53C21.

Key words and phrases. Convex integration, isotopy of knots, tantrix.

The first-named author was supported by NSF grant DMS-2202337.

Proof. After a reparametrization and, if necessary, an ambient homothety, we may assume that f has unit speed. Choose λ so large that the curvature of f is smaller than $\lambda\tilde{\kappa}$. Then, by Theorem 1.1, f is \mathcal{C}^1 -close and therefore isotopic to a unit-speed curve \tilde{f} with curvature $\lambda\tilde{\kappa}$. Finally, \tilde{f} is isotopic to $\lambda\tilde{f}$, which has curvature $\tilde{\kappa}$. \square

The above corollary generalizes [13, Cor. 1], where $\alpha = 2$ and the speed was not constant. See also [4–6] where this result was obtained for constant $\tilde{\kappa}$. The existence of \mathcal{C}^2 knots of constant curvature was first established by McAtee [10].

As in [4–6], the proof of Theorem 1.1 is based on a variant of convex integration [2, 3, 7, 12]. It hinges on the fact that when $|f'| = 1$, $\kappa = |T'|$. After reparametrizing f with unit speed, we deform T to a (longer) spherical curve \tilde{T} with speed $\tilde{\kappa}$, which we then integrate to obtain \tilde{f} . For \tilde{f} to be \mathcal{C}^1 -close to f , \tilde{T} should be \mathcal{C}^0 -close to T . Moreover, for \tilde{f} to be tangent to f at the prescribed points $p_i \in \Gamma$, \tilde{T} should have the same integral as T on every interval between successive p_i . These requirements will be met by combining basic convex geometry (Lemma 4.2) with mapping degree theory (Lemma 4.1). The latter is the novel feature of this work, which makes the arguments significantly shorter than those in [4].

2. PRELIMINARIES

We begin by recording some basic facts from [4] which are needed here.

2.1. Unit speed reparametrization. It suffices to establish Theorem 1.1 for unit-speed curves. To see this, note that for any $f \in \mathcal{C}^1(\Gamma, \mathbf{R}^n)$ and $\varphi \in \mathcal{C}^1(\Gamma, \Gamma)$, we have [4, Lem. 5.1]

$$|f \circ \varphi|_1 \leq |f|_1(1 + |\varphi|_1).$$

Suppose that Theorem 1.1 holds for unit-speed curves. Given $f \in \text{Imm}^\alpha(\Gamma, \mathbf{R}^n)$, choose $\lambda > 0$ such that $\text{length}(\lambda f) = b - a$. Then there exists a \mathcal{C}^α diffeomorphism $\varphi: \Gamma \rightarrow \Gamma$ such that $\lambda f \circ \varphi$ has unit speed. Let $\widetilde{\lambda f \circ \varphi}$ be the unit-speed curve with curvature $\tilde{\kappa} \circ \varphi / \lambda$ such that

$$|\widetilde{\lambda f \circ \varphi} - \lambda f \circ \varphi|_1 \leq \varepsilon \lambda / (1 + |\varphi^{-1}|_1),$$

and $\widetilde{\lambda f \circ \varphi}$ is tangent to $\lambda f \circ \varphi$ at $\varphi^{-1}(p_i)$, where p_i are the prescribed points. Set $\tilde{f} := \widetilde{\lambda f \circ \varphi} \circ \varphi^{-1} / \lambda$. Then \tilde{f} has curvature $\tilde{\kappa}$, and is tangent to f at p_i . Furthermore,

$$|\tilde{f} - f|_1 = |(\widetilde{\lambda f \circ \varphi} - \lambda f \circ \varphi) \circ \varphi^{-1}|_1 / \lambda \leq |\widetilde{\lambda f \circ \varphi} - \lambda f \circ \varphi|_1 (1 + |\varphi^{-1}|_1) / \lambda \leq \varepsilon,$$

as desired.

2.2. Average and center of mass. Let $I := [a, b]$ and $|I| := b - a$. For any curve $f \in \text{Imm}^{\alpha \geq 1}(I, \mathbf{R}^n)$, set $\text{length}(f) := \int_I |f'| du$. The *average* and *center of mass* of f are defined as

$$\text{ave}(f) := \frac{1}{|I|} \int_I f dt, \quad \text{and} \quad \text{cm}(f) := \frac{1}{\text{length}(f)} \int_I f |f'| dt.$$

In particular, $\text{cm}(f) = \text{ave}(f)$ when f has constant speed. Moreover, if $g: [c, d] \rightarrow \mathbf{R}^n$ is a *reparametrization* of f , i.e., there exists a diffeomorphism $\varphi: [c, d] \rightarrow [a, b]$ such

that $g = f \circ \varphi$, then $\text{cm}(f) = \text{cm}(g)$. For any positive (density) function $\rho \in \mathcal{C}^0(I, \mathbf{R})$, we define the corresponding *mass* and *center of mass* of f as

$$\text{m}(f, \rho) := \int_I \rho |f'| dt, \quad \text{and} \quad \text{cm}(f, \rho) := \frac{1}{\text{m}(f, \rho)} \int_I f \rho |f'| dt.$$

So $\text{cm}(f, 1/|f'|) = \text{ave}(f)$. More generally, letting $\varphi: [0, \text{m}(f, \rho)] \rightarrow I$ be the inverse of the (mass) function $t \mapsto \int_a^t \rho |f'| du$, we obtain

$$(1) \quad \text{cm}(f, \rho) = \text{ave}(f \circ \varphi),$$

see [4, Lem. 2.2].

3. REDUCTION TO SPHERICAL CURVES

Here we show that Theorem 1.1 follows from a geometric result (Proposition 3.2) for spherical curves. First we reduce Theorem 1.1 to the following local problem.

Proposition 3.1. *Let $f \in \text{Imm}^{\alpha \geq 2}(I, \mathbf{R}^n)$ be a unit-speed curve with curvature $\kappa > 0$, and V be an open neighborhood of $T(I)$ in \mathbf{S}^{n-1} . Then for any $\mathcal{C}^{\alpha-2}$ function $\tilde{\kappa}: I \rightarrow \mathbf{R}$ with $\tilde{\kappa} > \kappa$, there exists a unit-speed curve $\tilde{f} \in \text{Imm}^\alpha(I, \mathbf{R}^n)$ with curvature $\tilde{\kappa}$ such that $\tilde{T}(I) \subset V$, $f = \tilde{f}$ on ∂I , and $T(U) = \tilde{T}(\tilde{U})$ for some open neighborhoods U, \tilde{U} of ∂I in I .*

To see that the above proposition implies Theorem 1.1 recall that, as discussed in Section 2.1, we may assume that f in Theorem 1.1 has unit speed. So $T = f'$. Let I_i be a partition of Γ into intervals such that ∂I_i include all the prescribed points, and set $f_i := f|_{I_i}$. Choose I_i so small that $|I_i| \leq 1$ and $T(I_i)$ lies in the interior V_i of a ball of radius $\varepsilon/2$ in \mathbf{S}^{n-1} . Applying Proposition 3.1 to f_i , we obtain a \mathcal{C}^α curve \tilde{f}_i with unit speed and curvature $\tilde{\kappa}_i := \tilde{\kappa}|_{I_i}$ such that $\tilde{T}_i(I_i) \subset V_i$, $f_i = \tilde{f}_i$ on ∂I_i , and $T_i(U_i) = \tilde{T}_i(\tilde{U}_i)$ for some open neighborhoods U_i, \tilde{U}_i of ∂I_i in I_i .

Define \tilde{f} by $\tilde{f}|_{I_i} := \tilde{f}_i$. Then \tilde{f} is \mathcal{C}^0 , because $f_i = \tilde{f}_i$ on ∂I_i . Moreover \tilde{f} is \mathcal{C}^1 , because $\tilde{T}_i = \tilde{f}'_i$ and $\tilde{T}_i = T_i$ on ∂I_i . Hence $\tilde{T} := \tilde{f}'$ is well-defined. Note that \tilde{T} is piecewise $\mathcal{C}^{\alpha-1}$. Furthermore, since $T_i(U_i) = \tilde{T}_i(\tilde{U}_i)$, there are open neighborhoods W_i and \tilde{W}_i of ∂I_i in Γ such that $\tilde{T}(\tilde{W}_i) = T(W_i)$. Hence \tilde{T} is a reparametrization of T with speed $\tilde{\kappa}$ near ∂I_i , and so is $\mathcal{C}^{\alpha-1}$. Consequently \tilde{f} is \mathcal{C}^α and has curvature $\tilde{\kappa}$. Next note that for $t \in I_i$, $\tilde{f}(t) = f(t_i) + \int_{t_i}^t \tilde{T}_i du$ and $f(t) = f(t_i) + \int_{t_i}^t T_i du$, where t_i is the initial point of I_i . Furthermore, since $\tilde{T}(I_i) \subset V_i$, we have $|\tilde{T} - T| \leq \varepsilon$. Thus,

$$|\tilde{f}(t) - f(t)| = \left| \int_{t_i}^t (\tilde{T}_i - T_i) du \right| \leq \int_{t_i}^t |\tilde{T}_i - T_i| du \leq \varepsilon |I_i| \leq \varepsilon.$$

Also $|\tilde{f}' - f'| = |\tilde{T} - T| \leq \varepsilon$. Thus, after replacing ε by $\varepsilon/2$ if necessary, $|\tilde{f} - f|_1 \leq \varepsilon$. So Proposition 3.1 does indeed imply Theorem 1.1. Next we show that Proposition 3.1 is a consequence of the following result for spherical curves. We say that a curve $f \in \text{Imm}^\alpha(I, \mathbf{R}^n)$ is *nonflat* if the convex hull of $f(I)$ has interior points.

Proposition 3.2. *Let $T \in \text{Imm}^{\alpha \geq 1}(I, \mathbf{S}^{n-1})$ be a nonflat curve. Then for any $\mathcal{C}^{\alpha-1}$ function $\tilde{v}: I \rightarrow \mathbf{R}$ with $\tilde{v} > |T'|$ and open neighborhood V of $T(I)$, there exists a curve $\tilde{T} \in \text{Imm}^{\alpha}(I, \mathbf{S}^{n-1})$ such that $|\tilde{T}'| = \tilde{v}$, $\tilde{T}(I) \subset V$, $T(U) = \tilde{T}(\tilde{U})$ for some open neighborhoods U, \tilde{U} of ∂I , and $\text{ave}(T) = \text{ave}(\tilde{T})$.*

To see that Proposition 3.2 implies Proposition 3.1, let T be the tantrix of f in Proposition 3.1. After a small \mathcal{C}^{α} perturbation of f on a compact set in the interior of I , we may assume that T is nonflat. This is possible since by assumption $\kappa > 0$. Hence f does not trace a line segment, and so it can be perturbed, through unit-speed curves and with support in a compact subinterval of I , so that its tantrix becomes nonflat. Furthermore, since the perturbation is \mathcal{C}^2 -close to the original curve, we will still have $\tilde{\kappa} > \kappa$. Now let \tilde{T} be the $\mathcal{C}^{\alpha-1}$ curve obtained by applying Proposition 3.2 to T with $\tilde{v} := \tilde{\kappa}$. Then $\tilde{f}(t) := f(a) + \int_a^t \tilde{T} du$ is a \mathcal{C}^{α} curve with $\tilde{f}' = \tilde{T}$. So \tilde{f} has unit speed and curvature $\tilde{\kappa}$. Finally, the assumption that $\text{ave}(T) = \text{ave}(\tilde{T})$ ensures that $f = \tilde{f}$ on ∂I , which completes the argument.

4. PROOF OF THEOREM 1.1

It remains to establish Proposition 3.2, which completes the proof of Theorem 1.1 as discussed above. Our proof is based on the following topological lemma, which is established quickly via basic degree theory.

Lemma 4.1 ([6]). *Let $B \subset \mathbf{R}^n$ be a ball of radius R centered at x_0 , and $F: B \rightarrow \mathbf{R}^n$ be a continuous map. If $|F(x) - x| < R$ for all $x \in \partial B$, then $x_0 \in F(B)$.*

We also need the following version of a classical result of Kalman [8], who constructed continuous barycentric coordinates in convex polytopes.

Lemma 4.2. *Let $p_1, \dots, p_k \in \mathbf{R}^n$ be a collection of points whose convex hull has nonempty interior, and B be a ball centered at $x_0 := \sum_{i=1}^k p_i/k$. If B is sufficiently small, then there are positive \mathcal{C}^{∞} functions $c_i: B \rightarrow \mathbf{R}$ such that*

$$\sum_{i=1}^k c_i(x) = 1, \quad \sum_{i=1}^k c_i(x)p_i = x, \quad c_i(x_0) = \frac{1}{k}.$$

Proof. We may assume that $x_0 = 0$ after a translation. Then there are n linearly independent vectors among p_i , say p_1, \dots, p_n . So there are unique coefficients $a_1(x), \dots, a_n(x)$ such that $x = \sum_{i=1}^n a_i(x)p_i$ for each $x \in B$. The functions $a_i: B \rightarrow \mathbf{R}$ are \mathcal{C}^{∞} , as they are linear. Set $a_i = 0$ for $i > n$, and let

$$c_i(x) := \frac{1}{k} \left(1 - \sum_{j=1}^k a_j(x) \right) + a_i(x).$$

Since $\sum_{i=1}^n a_i(x_0)p_i = \sum_{i=1}^k a_i(x_0)p_i = x_0$ and p_1, \dots, p_n are linearly independent, $a_i(x_0) = 0$. Thus $c_i(x_0) = 1/k$. So $c_i > 0$ on B if B is small. Furthermore,

$$\begin{aligned} \sum_{i=1}^k c_i(x) &= 1 - \sum_{i=1}^k a_i(x) + \sum_{i=1}^k a_i(x) = 1, \\ \sum_{i=1}^k c_i(x)p_i &= x_0 \left(1 - \sum_{i=1}^k a_i(x) \right) + \sum_{i=1}^k a_i(x)p_i = x, \end{aligned}$$

as desired. \square

Now we are ready to prove the key result of this work:

Proof of Proposition 3.2. Let $\psi: [0, 1] \rightarrow I$ be given by $\psi(t) := (b-a)t + a$. Then $\text{ave}(T \circ \psi) = \text{ave}(T)$. Let $v := |T'|$. After replacing T with $T \circ \psi$, and v, \tilde{v} with $(b-a)v \circ \psi, (b-a)\tilde{v} \circ \psi$, respectively, we may assume that $I = [0, 1]$. Then

$$x_0 := \text{ave}(T) = \int_I T.$$

Since T is nonflat, there exists a ball B of radius $R > 0$ in the convex hull of $T(I)$ centered at x_0 . By Lemma 4.1 it is enough to construct a continuous mapping

$$B \ni x \mapsto \tilde{T}_x \in \text{Imm}^\alpha(I, \mathbf{S}^{n-1})$$

such that \tilde{T}_x satisfies the first three required properties in the statement of the proposition, and $|\text{ave}(\tilde{T}_x) - x| < R$. To this end we first construct a continuous family of reparametrizations $\bar{T}_x \in \text{Imm}^\alpha(I, \mathbf{S}^{n-1})$ of T such that

$$(2) \quad |\text{ave}(\bar{T}_x) - x| < R/2, \quad \text{and} \quad \bar{T}_{x_0} = T.$$

Then we construct \tilde{T}_x by adding certain loops to \bar{T}_x so that $|\text{ave}(\tilde{T}_x) - \text{ave}(\bar{T}_x)| < R/2$, which will complete the proof.

(Part I) We need to find a continuous family of \mathcal{C}^α diffeomorphisms $\varphi_x: I \rightarrow I$ such that $\bar{T}_x := T \circ \varphi_x$ satisfies the conditions listed in (2). To this end it suffices to construct a continuous family of positive $\mathcal{C}^{\alpha-1}$ functions $\rho_x: I \rightarrow \mathbf{R}$ such that

$$(3) \quad |\text{cm}(T, \rho_x) - x| < R/2, \quad \rho_{x_0} = 1/|T'|, \quad \text{and} \quad \int_I \rho_x |T'| dt = 1.$$

Then the inverse of the function $[0, 1] \ni t \mapsto \int_0^t \rho_x |T'| du$ gives the desired φ_x . Indeed the last condition in (3) ensures that $\varphi_x: I \rightarrow I$, and since ρ_x is $\mathcal{C}^{\alpha-1}$ and positive, φ_x is a \mathcal{C}^α diffeomorphism by the inverse function theorem. Furthermore, the first condition in (3) yields the first condition in (2) by (1). Finally, the second condition in (3) guarantees that φ_{x_0} is the identity, which yields the second condition in (2).

To construct ρ_x we rewrite (3) as

$$(4) \quad \left| \int_I \bar{\rho}_x T dt - x \right| < R/2, \quad \bar{\rho}_{x_0} = 1, \quad \text{and} \quad \int_I \bar{\rho}_x = 1,$$

where $\bar{\rho}_x := |T'| \rho_x$. Hence it suffices to construct a continuous family of positive $\mathcal{C}^{\alpha-1}$ functions $\bar{\rho}_x: I \rightarrow \mathbf{R}$ satisfying (4). To this end let I_i be a partition of I into $k > n$ equal segments, and θ_i be a \mathcal{C}^∞ partition of unity subordinate to I_i . Set

$$p_i := k \int_I \theta_i T dt.$$

Then

$$x_0 = \sum_i \int_I \theta_i T dt = \frac{1}{k} \sum_i p_i.$$

Assuming k is sufficiently large, the convex hull of p_i , $\text{conv}(\{p_i\})$, has interior points, since T is nonflat. In particular x_0 lies in the interior of $\text{conv}(\{p_i\})$. Fix k and choose R so small that B lies in $\text{conv}(\{p_i\})$. Then, by Lemma 4.2, there exist positive \mathcal{C}^∞ functions $c_i: B \rightarrow \mathbf{R}$ with $\sum c_i = 1$ such that $\sum c_i(x) p_i = x$ and $c_i(x_0) = 1/k$. Set

$$\bar{\rho}_x := \lambda(x) \sum_i c_i(x) \theta_i, \quad \text{where} \quad \lambda(x) := 1 / \int_I \sum_i c_i(x) \theta_i dt.$$

Then $\bar{\rho}_{x_0} = 1$, and $\int_I \bar{\rho}_x = 1$. Furthermore,

$$\int_I \bar{\rho}_x T dt = \lambda(x) \sum_i c_i(x) \int_I \theta_i T dt = \frac{\lambda(x)}{k} \sum_i c_i(x) p_i = \frac{\lambda(x)}{k} x.$$

We may assume that $x_0 = 0$ after a translation. Indeed, the quantity $\int_I \bar{\rho}_x T dt - x$ is unchanged under $T \mapsto T - x_0$, $x \mapsto x - x_0$, since $\int_I \bar{\rho}_x dt = 1$. Now, since $\lambda(x_0) = k$, choosing R sufficiently small we can make sure that $|\lambda(x)/k - 1| < 1/2$ for all $x \in B$. Then, since $|x| < R$,

$$\left| \int_I \bar{\rho}_x T dt - x \right| = \left| \frac{\lambda(x)}{k} - 1 \right| |x| < R/2,$$

as desired.

(Part II) Partition I into subintervals I_i such that for all $x \in B$, \bar{T}_x is injective on I_i and $\text{length}(\bar{T}_x|_{I_i}) < R/2$. So $\bar{T}_x(I_i)$ lies in a ball U_i^x of radius less than $R/4$ centered at the midpoint q_i^x of $\bar{T}_x(I_i)$. Let $C_i^x \subset (V \cap U_i^x)$ be a family of \mathcal{C}^α loops, depending continuously on q_i^x , that coincide with $\bar{T}_x(I_i)$ near q_i^x . For instance, we may construct C_i^x by gluing a segment of $\bar{T}_x(I_i)$ to an arc of a circle centered at q_i^x and rounding off the corners. Similarly we may construct a continuous family of \mathcal{C}^α loops $C_{i,\ell}^x$ nested inside C_i^x with length $0 < \ell \leq \text{length}(C_i^x)$ that contain a neighborhood of q_i^x in $\bar{T}_x(I_i)$; see Figure 1. Now for any $L > 0$ we can define a unique composite loop of length L at q_i^x as follows. Let m be the largest integer satisfying $m \text{length}(C_i^x) \leq L$. Then go m laps around C_i^x , in the direction induced by \bar{T}_x , and one lap around $C_{i,\ell}^x$ for $\ell = L - m \text{length}(C_i^x)$, if $\ell > 0$. Since $\tilde{v} > |T'| = |\bar{T}'_{x_0}|$, we may choose R so small that $\tilde{v} > |\bar{T}'_x|$ for all $x \in B$. Then

$$\ell_i^x := \int_{I_i} \tilde{v} dt - \text{length}(\bar{T}_x|_{I_i}) > 0.$$




FIGURE 1.

Finally, let \tilde{T}_x be the curve of speed \tilde{v} tracing T plus loops of length ℓ_i^x at q_i^x . Then $\bar{T}_x(I_i) \subset \tilde{T}_x(I_i) \subset U_i^x$. Thus $|\tilde{T}_x - \bar{T}_x|_0 < R/2$, which yields that $|\text{ave}(\tilde{T}_x) - \text{ave}(\bar{T}_x)| < R/2$, as desired. \square

ACKNOWLEDGMENTS

We thank Andrzej Świąch for suggesting the proof of Lemma 4.2, and Micha Wasem for comments on an earlier draft of this work.

REFERENCES

- [1] S. Conti, C. De Lellis, and L. Székelyhidi Jr., *h-Principle and rigidity for $C^{1,\alpha}$ isometric embeddings*, Nonlinear partial differential equations, 2012, pp. 83–116. MR3289360 [↑1](#)
- [2] Y. Eliashberg and N. Mishachev, *Introduction to the h-principle*, Graduate Studies in Mathematics, vol. 48, American Mathematical Society, Providence, 2002. MR1909245 [↑1, 2](#)
- [3] H. Geiges, *h-Principles and flexibility in geometry*, Mem. Amer. Math. Soc. **164** (2003), no. 779, viii+58. MR1982875 [↑2](#)
- [4] M. Ghomi, *h-Principles for curves and knots of constant curvature*, Geom. Dedicata **127** (2007), 19–35. MR2338512 [↑1, 2, 3](#)
- [5] M. Ghomi and M. Raffaelli, *Deformations of curves with constant curvature*, Preprint, arXiv:2410.06027 (2024). (<https://arxiv.org/abs/2410.06027>). [↑1, 2](#)
- [6] ———, *h-Principles for curves and knots of constant torsion*, Comment. Math. Helv., to appear (2025). (<https://arxiv.org/abs/2407.01729>). [↑1, 2, 4](#)
- [7] M. Gromov, *Partial differential relations*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 9, Springer, Berlin, 1986. MR864505 [↑1, 2](#)
- [8] J. A. Kalman, *Continuity and convexity of projections and barycentric coordinates in convex polyhedra*, Pacific J. Math. **11** (1961), 1017–1022. MR133732 [↑4](#)
- [9] N. H. Kuiper, *On C^1 -isometric imbeddings. I, II*, Nederl. Akad. Wetensch. Proc. Ser. A **58** (1955), 545–556, 683–689. MR75640 [↑1](#)
- [10] J. M. McAtee Ganatra, *Knots of constant curvature*, J. Knot Theory Ramifications **16** (2007), no. 4, 461–470. MR2327737 [↑2](#)
- [11] J. Nash, *C^1 isometric imbeddings*, Ann. of Math. (2) **60** (1954), 383–396. MR65993 [↑1](#)
- [12] D. Spring, *Convex integration theory: Solutions to the h-principle in geometry and topology*, Monographs in Mathematics, vol. 92, Birkhäuser, Basel, 1998. MR1488424 [↑2](#)
- [13] M. Wasem, *h-Principle for curves with prescribed curvature*, Geom. Dedicata **184** (2016), 135–142. MR3547785 [↑1, 2](#)

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332

Email address: `ghomi@math.gatech.edu`

URL: `https://ghomi.math.gatech.edu`

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332

Email address: `raffaelli@math.gatech.edu`

URL: `https://matteoraffaelli.com`